

3.3. Negations as premises

3.3.0. Overview

A second group of rules for negation reverses the roles of an affirmative sentence and its negation.

3.3.1. Indirect proof

The basic principles for negation describe its role as a premise only in *reductio* arguments but a *reductio* is always available as an argument of last resort.

3.3.2. Using lemmas to complete *reductios*

The role negative resources play will be to contradict other sentences; since what they contradict must often be introduced as a lemma, a use of lemmas is built into the rule for exploiting negative resources.

3.3.3. More examples

These new rules permit some new approaches to entailments that could be established using the last section's rule; but they also support some further entailments.

Glen Helman 28 Aug 2008

3.3.1. Indirect proof

The last section pursued consequences of the law for negation as a conclusion. The rules of this section will implement the other basic law for negation, the law for it as a premise:

$$\Gamma, \neg \varphi \Rightarrow \perp \text{ if and only if } \Gamma \Rightarrow \varphi$$

This says that a negation is $\neg \varphi$ inconsistent with a set Γ if and only if the sentence φ is entailed by that set.

There are two lessons we can learn from this law. First, the **only-if**-statement tells us that negative conclusions are not the only ones that can be established by way of *reductio* arguments. That is, an entailment $\Gamma \Rightarrow \varphi$ can be supported by the *reductio* $\Gamma, \neg \varphi \Rightarrow \perp$. The **if**-statement tells us in part that such an approach is safe, that the *reductio* is valid whenever the argument we wish to support by it is valid. But **if**-statement tells us more. Notice that φ is just the sort of resource that would enable us to complete a *reductio* that has $\neg \varphi$ as a premise. The **if**-claim above tells us that, if a *reductio* with $\neg \varphi$ as a premise can be completed at all, we would be able to validly conclude φ as a lemma—and that concluding it would not depend on using $\neg \varphi$ itself as a premise. This further lesson will provide the basis for exploiting negative resources, but its full application depends on the broader use of *reductio* arguments supported by the other two lessons, and that is what we will consider first.

Here is an example of this broader use. If we take **No one is home** to be the negation \neg **someone is home**, the law for negation as a premise says we can rest the validity of the left-hand argument below on the validity of the right-hand argument.

If no one was out, the car was in the driveway	If no one was out, the car was in the driveway
The car wasn't in the driveway	The car wasn't in the driveway
Someone was out	No one was out
	\perp

The right-hand argument depends in part on the logical properties of **if**; but, as far as negation is concerned, it depends on only the fact that a sentence and its negation are mutually exclusive.

The same basic fact gives us $\neg \neg \varphi, \neg \varphi \Rightarrow \perp$. If we apply the law for negation as a premise to this, we get the principle $\neg \neg \varphi \Rightarrow \varphi$. Moreover, the latter principle can be combined with the law for negation as a conclusion to establish the law for negation as a premise. So the further

logical properties of negation that are captured by the law for negation as a premise can be summarized in the principle that a double negation entails the corresponding positive claim.

This principle is one that was rejected by Brouwer in his intuitionistic mathematics. And one of his chief reasons for rejecting it was that it would allow us to draw a conclusion of the form **Something has the property P** when the corresponding claim **Nothing has the property P** was inconsistent with our premises—that is, just the sort of thing done in the example above. His concern with this is that it would enable us to conclude **Something has the property P** in cases where we were unable, even in principle, to provide an actual example of a thing with that property P. Brouwer did not object to such an argument in ordinary reasoning about the physical world (like the example above); but he held that, in reasoning concerning infinite mathematical structures, we were not reasoning about an independently existing realm of objects but instead about procedures for constructing abstract objects and that we had no business claiming the existence of such objects without having procedures enabling us to construct them. Brouwer’s concerns may not lead you to question the law for negation as a premise; but they highlight the indirectness of basing a positive conclusion on the fact that its denial is inconsistent with our premises. This aspect of these arguments is reflected in a common term for them, *indirect proofs*.

Although we will employ indirect proofs, we will need them for only a limited range of conclusions. We have other ways of planning for a goal that is a conjunction or a negation. We can simply close a gap whose goal is \top . And we will not adopt any rule to plan for the goal \perp of a *reductio argument*. At the moment, that leaves only unanalyzed components; and, until the last chapter, those are the only goals for which we will use indirect proofs. We have often closed gaps whose goals are atomic so, even for them, we know that indirect proof is not always necessary. However, it will serve us as a last resort.

In chapter 6, we will begin to analyze sentences into components that are not sentences, and we will still use indirect proof for goals that are analyzed in that way. In anticipation of this, we will use the term *atomic* for the kind of goals to which we will apply indirect proof; and we will refer to other sentences as *non-atomic*. Until chapter 6, any sentence we analyze will be a compound formed by applying a connective to one or more sentences, so, for the time being, the atomic sentences will be the unanalyzed sentences. \top and \perp count as non-atomic since identifying

them as logical constants counts as an analysis of their logical form. As a result, for the time being, the atomic sentences will be simple letters, and all other sentences will be non-atomic.

The rule implementing indirect proofs in derivations will be called *Indirect Proof* (IP). It takes the following form:





Fig. 3.3.1-1. Developing a derivation by planning for an atomic sentence at stage n .

Here is an example, which is related to the argument at the beginning of 3.2.2.



This example adds to the premise **Ann and Bill were not both home without the car being in the driveway** further premises telling us that each of Ann and Bill was home, and we conclude that the car was in the driveway. Although the initial premises and conclusion differ from those of the argument in 3.2.2, the *reductio* argument that is set up at stage 1 here has the same resources as the *reductio* set up at stage 3 in the derivation for the argument of 3.2.2 that was given at the end of 3.2.3.

The rule IP is not direct (in the sense used in 2.3.4) because it introduces a sentence more complex than the goal it plans for. It is, however, progressive. We will treat both atomic sentences and their negations as equally basic when they are resources: neither sort of resource will be exploited. And, as was noted above, we will treat \perp as

the basic form of goal, the only one without a corresponding planning rule. Thus IP leaves us with a goal that requires no planning and introduces no resources that need to be exploited further. This is, of course, not to say that applying IP will eliminate the need for further exploitations; indeed, since negated compounds will be exploited only in *reductio* arguments, we will often be in a position to exploit such resources only after we have used IP. The rule we will use to do that is the one we will consider next.

Glen Helman 28 Aug 2008

3.3.2. Using lemmas to complete *reductios*

Now that we have IP, we are in a position to provide a proof for any argument whose validity depends only on the properties of \top , \perp , conjunction, and negation. However, to do this using only the rules we have so far, we would often need to use LFR—or, in simpler cases, Adj—to make use of negative resources. This poses no problems when we construct derivations for valid arguments, but it makes it difficult to show that any argument is not valid. LFR does not itself exploit resources, so negated compounds remain as active resources until a gap is closed. In order to count an open gap as having reached a dead end, we would need some description of the conditions under which LFR had been used often enough. Such a description could certainly be given; and, in the last two chapters, we will need to take an analogous approach in the case of one of the rules for quantifiers. But, in the case of negation, it is possible to keep track of the use of resources by way of a genuine exploitation rule.

The basis for such an approach was the third lesson drawn from the law of negation as a premise: if a *reductio* that has $\neg \phi$ as a premise is valid—that is, if $\Gamma, \neg \phi \Rightarrow \perp$ —then ϕ is a valid conclusion from the premises other than $\neg \phi$ (i.e., $\Gamma \Rightarrow \phi$). And ϕ is just the lemma we need in order to use the premise $\neg \phi$ to complete the *reductio*. That means not only that it is safe to introduce ϕ as a lemma but also that the gap in which we establish the lemma need not contain $\neg \phi$ among its active resources. Of course, $\neg \phi$ is needed along with the lemma to reach the goal \perp , but there is no need to introduce a second gap in which we try to reach this goal (as would be done with LFR) because such a gap would close immediately by Nc. And, indeed, the law for negation as a premise tells us not only that we can reach the needed lemma (provided that the *reductio* is valid to begin with) but also that reaching this lemma is all that we need to do, for it says both that the *reductio* argument $\Gamma, \neg \phi / \perp$ is valid only in cases when the argument Γ / ϕ is valid and that $\Gamma, \neg \phi / \perp$ is valid in all such cases.

We will call the rule that implements these ideas *Completing a Reductio* (CR).





Fig. 3.3.2-1. Developing a derivation by exploiting a negated compound at stage n .

The motivation for this rule lies in its use with the negations of non-atomic sentences; and, in fact, we must limit its use to such sentences. It is sound and safe in the case of atomic sentence, but it would not be progressive in that case (given the way we are measuring distance from a dead end) because it would replace a resource that we never exploit by a goal that we could go on to plan for by IP; that is, it would provide new opportunities for developing a derivation and thus send us farther from reaching a dead end. Both IP and CR carry us between gaps whose proximate arguments have the forms $\Gamma, \neg \varphi / \perp$ and Γ / φ ; but they carry us in opposite directions, so, if there is any overlap in the sentences φ to which they apply, a derivation could move back and forth between the two arguments forever. We block such circles by limiting IP to cases where φ is atomic and limiting CR to cases where φ is non-atomic.

The following derivations show, on the left, the use of CR in a derivation for the argument from 3.2.2 that was used as an illustration in the last subsection and, on the right, an analogous use of LFR:



The most obvious difference between the two is an extra argument in the second in which the lemma is $(A \wedge B) \wedge \neg C$ is used explicitly. But the more important difference is that, while the first premise is exploited at stage 2 in the left-hand derivation, it remains unexploited in the second.

These derivations are different in length, and the left-hand derivation is a little longer than the one at the end of 3.3.1. The length of a derivation is not important in its own right, but the differences in length reflect differences in the way these derivations are constructed. The extra length of these two in comparison with the earlier derivation is the result of introducing $(A \wedge B) \wedge \neg C$ as an explicit goal. The derivation above using CR is shorter than the one using LFR because CR applies only in cases where we know exactly how to use the lemma to complete the *reductio*. It is longer than the earlier derivation using just Adj because it provides us with a stage at which we can mark its negation as exploited and makes explicit the resource aimed at by the two uses of Adj.

3.3.s. Summary

- 1 The law for negation as a premise tells us two things about entailment. The first is that any conclusion is valid if and only if the denial of that conclusion can be reduced to absurdity given the premises. This is the principle of indirect proof; it is closely tied to the entailment $\neg \neg \varphi \Rightarrow \varphi$ (and is subject to the same concerns as is that entailment). We have no need for this principle except in the case of unanalyzed components, which we will begin to call atomic sentences. And, for reasons noted later, we need to limit the use of the rule Indirect Proof (IP) to them.
- 2 Another lesson we can draw from the law for negation as a premise is that a *reductio* argument with a negative premise $\neg \varphi$ is valid if and only if the sentence φ is entailed by the other premises. This tells us that φ can be safely introduced as a lemma even if we drop $\neg \varphi$ from our active resources. The rule implementing this idea, Completing a Reductio (CR) serves as our rule for exploiting negative resources. It applies only to *reductio* arguments but the availability of IP insures that any gap will eventually turn into a gap in a *reductio* argument (unless it closes before that point). Since CR, by dropping a resource $\neg \varphi$ and adding a goal φ has an effect opposite to that of IP, we must apply them to different sentences φ to avoid going in circles. So, just as IP is limited to atomic sentences, CR is limited to negations of non-atomic sentences.
- 3 The rule CR can lead us to set as goals any lemmas we need to use negations in completing *reductio* arguments. It therefore eliminates any need for LFR. The rule Adj is also no longer needed since the rules CR and Cnj will lead us to identify and prove any lemma that Adj would introduce. Indeed, derivations for arguments involving conjunction can now be constructed by letting the rules guide us completely.

Glen Helman 28 Aug 2008

3.3.x. Exercise questions

Use derivations to establish each of the claims of entailment shown below. You can maximize your practice in the use of CR by avoiding LFR and using Adj only when the goal is a conjunction.

1. $\neg (A \wedge \neg B), A \Rightarrow B$
2. $J \wedge \neg (J \wedge \neg C) \Rightarrow J \wedge C$ (see exercise 1j of 3.1.x)
3. $\neg (\neg (A \wedge B) \wedge C), \neg A \Rightarrow \neg C$
4. $\neg (A \wedge \neg (B \wedge C)) \Rightarrow \neg (A \wedge \neg B)$
5. $\neg (A \wedge \neg B), \neg (B \wedge \neg C) \Rightarrow \neg (A \wedge \neg C)$
6. $\neg (A \wedge \neg B), \neg (A \wedge \neg C) \Rightarrow \neg (A \wedge \neg (B \wedge C))$

For more exercises, use the exercise machine.

Glen Helman 28 Aug 2008

3.3.xa. Exercise answers

1.	$\neg(A \wedge \neg B)$ 2 A (3)
3 Adj	$\neg B$ (3) $A \wedge \neg B$ X,(4)
4 QED	$A \wedge \neg B$ 2
2 CR	\perp 1
1 IP	B
2.	$J \wedge \neg(J \wedge \neg C)$ 1
1 Ext	J (3),(6)
1 Ext	$\neg(J \wedge \neg C)$ 5
3 QED	J 2
6 Adj	$\neg C$ (6) $J \wedge \neg C$ X,(7)
7 QED	$J \wedge \neg C$ 5
5 CR	\perp 4
4 IP	C 2
2 Cnj	$J \wedge C$

3.	$\neg(\neg(A \wedge B) \wedge C)$ 2 $\neg A$ (7)
6 Ext	C (4)
6 Ext	$A \wedge B$ 6 A (7) B
7 Nc	\perp 5
5 RAA	$\neg(A \wedge B)$ 3 \bullet
4 QED	C 3
3 Cnj	$\neg(A \wedge B) \wedge C$ 2
2 CR	\perp 1
1 RAA	$\neg C$
4.	$\neg(A \wedge \neg(B \wedge C))$ 3
2 Ext	$A \wedge \neg B$ 2
2 Ext	A (5) $\neg B$ (8)
5 QED	A 4 \bullet
7 Ext	$B \wedge C$ 7
7 Ext	B (8) C
8 Nc	\perp 6
6 RAA	$\neg(B \wedge C)$ 4 \bullet
4 Cnj	$A \wedge \neg(B \wedge C)$ 3
3 CR	\perp 1
1 RAA	$\neg(A \wedge \neg B)$

	$\neg(A \wedge \neg B)$	3										
	$\neg(B \wedge \neg C)$	7										
	$A \wedge \neg C$	2										
2 Ext	A	(5)										
2 Ext	$\neg C$	(8)										
	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">A</td> </tr> </table>	•			A	4						
•												
	A											
5 QED	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">B</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge \neg C$</td> <td style="padding-left: 10px;">(8)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge \neg C$</td> </tr> </table> </td> </tr> </table>		B		<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge \neg C$</td> <td style="padding-left: 10px;">(8)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge \neg C$</td> </tr> </table>	$B \wedge \neg C$	(8)	•			$B \wedge \neg C$	X,(9)
	B											
	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge \neg C$</td> <td style="padding-left: 10px;">(8)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge \neg C$</td> </tr> </table>	$B \wedge \neg C$	(8)	•			$B \wedge \neg C$					
$B \wedge \neg C$	(8)											
•												
	$B \wedge \neg C$											
8 Adj	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">\perp</td> </tr> </table>		\perp	6								
	\perp											
9 QED	$\neg B$	4										
7 CR	$A \wedge \neg B$	3										
6 RAA	\perp	1										
4 Cnj	$\neg(A \wedge \neg C)$											
3 CR	\perp	1										
1 RAA	$\neg(A \wedge \neg C)$											

	$\neg(A \wedge \neg B)$	3										
	$\neg(A \wedge \neg C)$	7										
	$A \wedge \neg(B \wedge C)$	2										
2 Ext	A	(5),(9)										
2 Ext	$\neg(B \wedge C)$	10										
	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">A</td> </tr> </table>	•			A	4						
•												
	A											
5 QED	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">B</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">A</td> </tr> </table> </td> </tr> </table>		B		<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">A</td> </tr> </table>	•			A	(11)		
	B											
	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">A</td> </tr> </table>	•			A							
•												
	A											
9 QED	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">C</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge C$</td> <td style="padding-left: 10px;">(11)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge C$</td> </tr> </table> </td> </tr> </table>		C		<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge C$</td> <td style="padding-left: 10px;">(11)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge C$</td> </tr> </table>	$B \wedge C$	(11)	•			$B \wedge C$	X,(12)
	C											
	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">$B \wedge C$</td> <td style="padding-left: 10px;">(11)</td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;">•</td> <td></td> </tr> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">$B \wedge C$</td> </tr> </table>	$B \wedge C$	(11)	•			$B \wedge C$					
$B \wedge C$	(11)											
•												
	$B \wedge C$											
11 Adj	<table style="border-collapse: collapse; margin-left: 5px;"> <tr> <td style="border-left: 1px solid black; padding-left: 5px;"> </td> <td style="padding-left: 5px;">\perp</td> </tr> </table>		\perp	9								
	\perp											
12 QED	$\neg C$	8										
10 CR	$A \wedge \neg C$	7										
9 RAA	\perp	6										
8 Cnj	$\neg B$	4										
7 CR	$A \wedge \neg B$	3										
6 RAA	\perp	1										
4 Cnj	$\neg(A \wedge \neg(B \wedge C))$											
3 CR	\perp	1										
1 RAA	$\neg(A \wedge \neg(B \wedge C))$											

Choosing $\neg(B \wedge C)$ as the resource to exploit by CR at stage 3 would lead to a somewhat shorter and simpler derivation.