

# Appendices

## Appendix A. Reference

### A.0. Overview

#### A.1. Basic concepts

Definitions of entailment and related ideas

#### A.2. Logical forms

Forms expressed using one or more logical constants together with symbolic and English notation or readings

#### A.3. Truth tables

Tables that stipulate the meaning of the constants of truth-functional logic

#### A.4. Derivation rules

A guide to the use of derivation rules with links to the rules themselves

Glen Helman 05 Nov 2011

### A.1. Basic concepts

<i>Concept</i>	<i>Negative definition</i>	<i>Positive definition</i>
$\varphi$ is <i>entailed</i> by $\Gamma$ : $\Gamma \models \varphi$	There is no logically possible world in which $\varphi$ is false while all members of $\Gamma$ are true.	$\varphi$ is true in every logically possible world in which all members of $\Gamma$ are true.
$\varphi$ and $\psi$ are ( <i>logically equivalent</i> ): $\varphi \simeq \psi$	There is no logically possible world in which $\varphi$ and $\psi$ have different truth values.	$\varphi$ and $\psi$ have the same truth value as each other in every logically possible world.
$\varphi$ is a <i>tautology</i> : $\models \varphi$ (or $\top \models \varphi$ )	There is no logically possible world in which $\varphi$ is false.	$\varphi$ is true in every logically possible world.
$\varphi$ is <i>inconsistent with</i> $\Gamma$ : $\Gamma, \varphi \models$ (or $\Gamma, \varphi \models \perp$ )	There is no logically possible world in which $\varphi$ is true while all members of $\Gamma$ are true.	$\varphi$ is false in every logically possible world in which all members of $\Gamma$ are true.
$\Gamma$ is <i>inconsistent</i> : $\Gamma \models$ (or $\Gamma \models \perp$ )	There is no logically possible world in which all members of $\Gamma$ are true.	In every logically possible world, at least one member of $\Gamma$ is false.
$\varphi$ is <i>absurd</i> : $\varphi \models$ (or $\varphi \models \perp$ )	There is no logically possible world in which $\varphi$ is true.	$\varphi$ is false in every logically possible world.
$\Sigma$ is <i>rendered exhaustive</i> by $\Gamma$ : $\Gamma \models \Sigma$	There is no logically possible world in which all members of $\Sigma$ are false while all members of $\Gamma$ are true.	At least one member of $\Sigma$ is true in each logically possible world in which all members of $\Gamma$ are true.

Glen Helman 05 Nov 2011

## A.2. Logical forms

### Forms for which there is symbolic notation

	Symbolic notation	English notation or English reading
Negation	$\neg \phi$	not $\phi$
Conjunction	$\phi \wedge \psi$	both $\phi$ and $\psi$ ( $\phi$ and $\psi$ )
Disjunction	$\phi \vee \psi$	either $\phi$ or $\psi$ ( $\phi$ or $\psi$ )
The conditional	$\phi \rightarrow \psi$ $\psi \leftarrow \phi$	if $\phi$ then $\psi$ ( $\phi$ implies $\psi$ ) yes $\psi$ if $\phi$ ( $\psi$ if $\phi$ )
Identity	$\tau = \upsilon$	$\tau$ is $\upsilon$
Predication	$\theta \tau_1 \dots \tau_n$	$\theta$ fits $\tau_1, \dots, \tau_n$ A series of terms $\tau_1, \dots, \tau_n$ can be read (series) $\tau_1, \dots$ , <b>an</b> $\tau_n$ (using the expression <b>an</b> to distinguish this use of <b>and</b> from its use in conjunction and adding <b>series</b> when necessary to avoid ambiguity)
Compound term	$\gamma \tau_1 \dots \tau_n$	$\gamma$ of $\tau_1, \dots, \tau_n$ $\gamma$ applied to $\tau_1, \dots, \tau_n$
Predicate abstract	$[\phi]_{x_1 \dots x_n}$	what $\phi$ says of $x_1 \dots x_n$
Functor abstract	$[\tau]_{x_1 \dots x_n}$	$\tau$ for $x_1 \dots x_n$
Universal quantification	$\forall x \theta x$	forall $x$ $\theta x$ everything, $x$ , is such that $\theta x$
Restricted universal	$(\forall x: \rho x) \theta x$	forall $x$ st $\rho x$ $\theta x$ everything, $x$ , such that $\rho x$ is such that $\theta x$
Existential quantification	$\exists x \theta x$	forsome $x$ $\theta x$ something, $x$ , is such that $\theta x$
Restricted existential	$(\exists x: \rho x) \theta x$	forsome $x$ st $\rho x$ $\theta x$ something, $x$ , such that $\rho x$ is such that $\theta x$
Definite description	$!x \rho x$	the $x$ st $\rho x$ the thing, $x$ , such that $\rho x$

### Some paraphrases of other forms

#### Truth-functional compounds

neither $\phi$ nor $\psi$	$\neg (\phi \vee \psi)$
$\psi$ only if $\phi$	$\neg \phi \wedge \neg \psi$
$\psi$ unless $\phi$	$\neg \psi \leftarrow \neg \phi$

#### Generalizations

All Cs are such that ( ... they ... )	$(\forall x: x \text{ is a } C) \dots x \dots$
No Cs are such that ( ... they ... )	$(\forall x: x \text{ is a } C) \neg \dots x \dots$
Only Cs are such that ( ... they ... )	$(\forall x: \neg x \text{ is a } C) \neg \dots x \dots$
with: among Bs	add to the restriction: $x \text{ is a } B$
except Es	$\neg x \text{ is an } E$
other than $\tau$	$\neg x = \tau$

#### Numerical quantifier phrases

At least 1 C is such that ( ... it ... )	$(\exists x: x \text{ is a } C) \dots x \dots$
At least 2 Cs are such that ( ... they ... )	$(\exists x: x \text{ is a } C) (\exists y: y \text{ is a } C \wedge \neg y = x) (\dots x \dots \wedge \dots y \dots)$
Exactly 1 C is such that ( ... it ... )	$(\exists x: x \text{ is a } C) (\dots x \dots \wedge (\forall y: y \text{ is a } C \wedge \neg y = x) \neg \dots y \dots)$ or $(\exists x: x \text{ is a } C) (\dots x \dots \wedge (\forall y: y \text{ is a } C \wedge \dots y \dots) x = y)$

#### Definite descriptions (on Russell's analysis)

The C is such that ( ... it ... )	$(\exists x: x \text{ is a } C \wedge (\forall y: \neg y = x) \neg y \text{ is a } C) \dots x \dots$ or $(\exists x: x \text{ is a } C \wedge (\forall y: y \text{ is a } C) x = y) \dots x \dots$
-----------------------------------	--

Glen Helman 05 Nov 2011

## A.4. Derivation rules

### Basic system

Rules for developing gaps		
logical form	as a resource	as a goal
atomic sentence		IP
negation $\neg \phi$ (if $\phi$ not atomic & goal is $\perp$ )	CR	RAA
conjunction $\phi \wedge \psi$	Ext	Cnj
disjunction $\phi \vee \psi$	PC	PE
conditional $\phi \rightarrow \psi$ (if goal is $\perp$ )	RC	CP
universal $\forall x \theta x$	UI	UG
existential $\exists x \theta x$	PCh	NcP

In addition, if the conditions for applying a rule are met except for differences between co-aliases, then the rule can be applied and is notated by adding "="; QED= and Nc= are examples of this.

Rules for closing gaps			
when to close	resources		rule
co-aliases	resources	goal	
	$\phi$	$\phi$	QED
$\phi$ and $\neg \phi$	$\perp$		Nc
		$\top$	ENV
	$\perp$		EFQ
$\tau \multimap \upsilon$		$\tau = \upsilon$	EC
$\tau \multimap \upsilon$	$\neg \tau = \upsilon$	$\perp$	DC
$\tau_1 \multimap \upsilon_1, \dots, \tau_n \multimap \upsilon_n$	$P\tau_1 \dots \tau_n$	$P\upsilon_1 \dots \upsilon_n$	QED=
$\tau_1 \multimap \upsilon_1, \dots, \tau_n \multimap \upsilon_n$	$P\tau_1 \dots \tau_n$	$\perp$	Nc=

#### Detachment rules (optional)

required resources	rule
main	auxiliary
$\phi \rightarrow \psi$	$\frac{\phi}{\neg^\pm \psi}$ MPP
$\phi \vee \psi$	$\frac{\neg^\pm \phi \text{ or } \neg^\pm \psi}{\phi \text{ or } \psi}$ MTP
$\neg(\phi \wedge \psi)$	$\frac{\neg^\pm \phi \text{ or } \neg^\pm \psi}{\phi \text{ or } \psi}$ MPT

### Additional rules (not guaranteed to be progressive)


Attachment rules	
added resource	rule
$\phi \wedge \psi$	Adj
$\phi \rightarrow \psi$	Wk
$\phi \vee \psi$	Wk
$\neg(\phi \wedge \psi)$	Wk
$\tau = \upsilon$	CE
$\theta \upsilon_1 \dots \upsilon_n$	Cng
$\exists x \theta x$	EG

Rule for lemmas	
prerequisite	rule
the goal is $\perp$	LFR


## Diagrams

### Rules from chapter 2


#### Extraction (Ext)




#### Conjunction (Cnj)




#### Quod Erat Demonstrandum (QED)




*Ex Nihilo Verum* (ENV)




*Ex Falso Quodlibet* (EFQ)



Adjunction (Adj)




Lemma for *Reductio* (LFR)




Rules from chapter 3


Indirect Proof (IP)




Completing the *Reductio* (CR)



*Reductio ad Absurdum (RAA)*




Non-contradiction (Nc)




Rules from chapter 4


Proof by Cases (PC)



*Proof of Exhaustion (PE)*



*Modus Tollendo Ponens (MTP)*



Modus Ponendo Tollens (MPT)

$$\frac{\begin{array}{c} \varphi \text{ [available]} \\ \dots \\ \neg(\varphi \wedge \psi) \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \varphi \quad (n) \\ \dots \\ \neg(\varphi \wedge \psi) \quad n \\ \dots \\ \vdots \\ \vdots \\ \hline \neg^{\pm} \psi \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ MPT}$$

$$\frac{\begin{array}{c} \psi \text{ [available]} \\ \dots \\ \neg(\varphi \wedge \psi) \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \psi \quad (n) \\ \dots \\ \neg(\varphi \wedge \psi) \quad n \\ \dots \\ \vdots \\ \vdots \\ \hline \neg^{\pm} \varphi \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ MPT}$$

Weakening (Wk)

$$\frac{\begin{array}{c} \dots \\ \varphi \text{ [available]} \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \dots \\ \varphi \quad (n) \\ \dots \\ \vdots \\ \vdots \\ \hline \varphi \vee \psi \quad X \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ Wk}$$

$$\frac{\begin{array}{c} \dots \\ \psi \text{ [available]} \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \dots \\ \psi \quad (n) \\ \dots \\ \vdots \\ \vdots \\ \hline \varphi \vee \psi \quad X \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ Wk}$$

Weakening (Wk)

$$\frac{\begin{array}{c} \dots \\ \neg^{\pm} \varphi \text{ [available]} \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \dots \\ \neg^{\pm} \varphi \quad (n) \\ \dots \\ \vdots \\ \vdots \\ \hline \neg(\varphi \wedge \psi) \quad X \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ Wk}$$


$$\frac{\begin{array}{c} \dots \\ \neg^{\pm} \psi \text{ [available]} \\ \dots \\ \vdots \\ \vdots \\ \hline \theta \\ \dots \end{array}}{\begin{array}{c} \dots \\ \neg^{\pm} \psi \quad (n) \\ \dots \\ \vdots \\ \vdots \\ \hline \neg(\varphi \wedge \psi) \quad X \\ \dots \\ \hline \theta \\ \dots \end{array}} \quad n \text{ Wk}$$

Rules from chapter 5


Rejecting a Conditional (RC)

$$\frac{\begin{array}{c} \dots \\ \varphi \rightarrow \psi \\ \dots \\ \vdots \\ \vdots \\ \hline \perp \\ \dots \end{array}}{\begin{array}{c} \dots \\ \varphi \rightarrow \psi \quad n \\ \dots \\ \vdots \\ \vdots \\ \hline \varphi \quad n \\ \vdots \\ \hline \psi \\ \vdots \\ \hline \perp \quad n \\ \dots \\ \hline \perp \\ \dots \end{array}} \quad n \text{ RC}$$


Conditional Proof (CP)




Modus Ponendo Ponens (MPP)




Modus Tollendo Tollens (MTT)



Weakening (Wk)




Weakening (Wk)




Rules from chapter 6


Equated Co-aliases (EC)



Distinguished Co-aliases (DC)




QED given equations (QED=)




Note: Two series of terms are co-alias series when their corresponding members are co-aliases.

Non-contradiction given equations (Nc=)




Note: Two series of terms are co-alias series when their corresponding members are co-aliases.

Co-alias Equation (CE)




Congruence (Cng)




Note:  $\theta$  can be an abstract, so  $\theta\tau_1 \dots \tau_n$  and  $\theta v_1 \dots v_n$  are any formulas that differ only in the occurrence of terms and in which the corresponding terms are co-aliases.

Rules from chapter 7

Universal Instantiation (UI)




Universal Generalization (UG)






Proof by Choice (PCh)



Non-constructive Proof (NcP)



Existential Generalization (EG)

